

PARCHMENT SKIN

78-year old male patient with parchment skin and a traumatic wound after falling

PREVIOUS TREATMENT

The treatment with ① started directly after the fall.

TREATMENT WITH ①

① was applied onto the wound and surrounding skin area after cleansing with water. The wound was covered with a fatty gauze for 10 days and with a non-woven gauze held in place with a self-adhesive gauze bandage for the remaining treatment.

RESULT

Wound closure within 21 days.


CONCLUSION HEALTH CARE PROFESSIONAL

Karin Eggenberger, wound expert, Switzerland

“Despite parchment skin trauma being a simple acute wound, its treatment is often challenging. The main problem is the potential adherence of a secondary dressing, which can lead to a traumatic wound opening when changing the dressing. The application of ① in combination with a fatty gauze prevents this. It was the patient's wish to use a fatty gauze as secondary dressing as he was very scared of pain during the dressing change. If applied daily, a non-woven gauze as secondary dressing is enough, as the oil film prevents adherence of the gauze as well. In addition, the peri-wound skin is nurtured when using ① and the newly formed skin seems to be very robust.”



Day 1



Day 7



Day 14



Day 21

MANUFACTURER

Phytoceuticals AG
CH-8134 Adliswil

FURTHER INFORMATION

www.wound.ch
Tel. +41 58 800 58 58

